

## Cotoneaster

## **Species Description**

Scientific name: Cotoneaster species

**AKA:** Cotoneasterau (Welsh)

Native to: Majority of species originate from east Asia

Habitat: Rough ground

A large group of small trees and prostrate shrubs that can be either evergreen or deciduous. They are becoming increasingly naturalised due to birds which eat the small red berries and spread the seed. There is one native species, wild Cotoneaster (*Cotoneaster cambricus*) which occurs as a few individuals in North Wales. This ID sheet includes those introduced species which are relatively common. Where they become established they can become dominant to the exclusion of native species.

Himalayan cotoneaster (*Cotoneaster simonsii*) is an erect deciduous shrub 3-4 metres high with 1.5-2.5 cm long leaves; small-leaved cotoneaster (*Cotoneaster microphyllou*) is an evergreen low-growing shrub with very small leaves (0.5-0.8cm long) and tree cotoneaster (*Cotoneaster* 

*frigidus*) is a deciduous or semi-evergreen shrub or small tree with leaves between 6-15 cm long and flowers in dense clusters. All these species have leaves which are shiny and hairless on the upper surface and slightly hairy on the lower surface. Cotoneaster species do not have thorns.

Wall cotoneaster (*Cotoneaster horizontalis*) is the most widely recorded species and is distinctive in having stems that spread horizontally in flattened herring-bone like branches and bears single flowers. Unlike the other cotoneasters mentioned, the underside of leaves of this species are relatively hairless.

Cotoneasters have been in cultivation in GB since 1824 and there are over 100 species now widely cultivated in the UK. Other species could also become naturalised.


Cotoneaster species are listed under Schedule 9 to the Wildlife and Countryside Act 1981 with respect to England and Wales. As such, it is an offence to plant or otherwise cause these species to grow in the wild.


For details of legislation go to www.nonnativespecies.org/legislation.



Himalayan cotoneaster (Cotoneaster simonsii)

## **Key ID Features**








## Identification throughout the year

Evergreen species such as the small-leaved cotoneaster can be identified throughout the year by its leaves. Deciduous species can be identified by the presence of the red berries which can remain throughout winter and, in the case of wall cotoneaster, by the distinctive herringboneshaped branches.









Other similar species which may be confused with cotoneaster include:

**Escallonia** (*Escallonia macrantha*) - an evergreen shrub which has alternate serrated leaves and numerous pinkish-red flowers, no thorns;

**Barberry (Berberis vulgaris)** - a thorny shrub with small serrated leaves, yellow flowers and red lozenge-shaped berries; and **Sea buckthorn (Hippophae rhamnoides)** - a thorny densely branched shrub with alternate long thin leaves with bright orange berries on female plants.





References and further reading:

Johnson, O and More, D (2004) "Collins Tree Guide". HarperCollins

Preston, C D, Pearman, D A and Dines, T A (editors) (2002) "New Atlas of the British and Irish Flora". Oxford University Press

Stace, C (1999) "Field Flora of the British Isles". Cambridge University Press